

# Erratum to: Co-culture of Retinal and Endothelial Cells Results in the Modulation of Genes Critical to Retinal Neovascularization

Ravindra Kumar, <sup>1, a, \*</sup> Sandra Harris-Hooker, <sup>2</sup> Ritesh Kumar, <sup>3</sup> Gary Sanford, <sup>1</sup>  
 @ corresponding author, & equal contributor

Vascular Cell. 2012; **4**(1):6 | © Kumar et al  
 Received: 14 March 2012 | Accepted: 26 March 2012 | Published: 26 March 2012  
 Vascular Cell ISSN: 2045-824X  
 DOI: <https://doi.org/10.1186/2045-824X-4-6>

### Author information

1. Department of Microbiology, Biochemistry and Immunology - Morehouse School of Medicine; Atlanta, 30310, Georgia, USA
2. Department of Pathology - Morehouse School of Medicine; Atlanta, 30310, Georgia, USA
3. Undergraduate student, Department of Chemistry and Biochemistry - Georgia Institute of Technology; Atlanta, 30322, Georgia, USA

[a] [rkumar@msm.edu](mailto:rkumar@msm.edu)

### Correction

Following publication of our article [1] it was noted that Figures five E and five G were the same as Figures six A and six B. Figure 1 in this correction

article is the correct version of Figure six that should have been included in the original article [1]. We apologize for any inconvenience caused by this error.




Figure 1

### Figure 1 caption

RT-PCR and Western blot analysis of NFκB. Total RNA and total protein were extracted from HRPC and HUVEC cultured alone or co-cultured under normoxia and hypoxia conditioned for 24 h. The expression of NFκB was measured by (A) electrophoresis of RT-PCR, (C) Western blot analysis in the HRPC and HUVEC. Figures (B, D) the band intensities corresponding to the NFκB were quantified by image analysis using a Bio-Rad scanning densitometer and Quantity One analysis software. Data were shown as ratio of NFκB densities to that of 18S RNA for RT-PCR and β-actin antibody was used to normalize Western blot for differences in loading and the transfer efficiencies. All data were expressed as mean +/- SE and results are representatives of three independent experiments.

## Authors' original submitted files for images

Authors' original file for figure 1  
[Click here to view.](#)

Below are the links to the authors' original submitted files for images.

## References

1. Kumar R, Harris-Hooker S, Kumar R, Sanford G. Co-culture of Retinal and Endothelial Cells Results in the Modulation of Genes Critical to Retinal Neovascularization. *Vascular Cell*. 2011;3:27-.

## Copyright & License

*Statement:* Copyright © 2012, Kumar et al.

*Holder:* Kumar et al

*Licensee:* Publiverse Online S.R.L.

*License: Open Access* This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.



The present article has been published in Vascular Cell journal by Publiverse Online S.R.L.